

Risk of Burnout in Perioperative Clinicians

A Survey Study and Literature Review

Steve A. Hyman, M.D.,* Damon R. Michaels, B.S.,† James M. Berry, M.D.,‡
Jonathan S. Schildcrout, Ph.D.,§ Nathaniel D. Mercaldo, M.S.,|| Matthew B. Weinger, M.D.#

ABSTRACT

Background: Burnout can lead to health and psychologic problems and is apparently increasing in physicians and nurses. Previous studies have not evaluated all healthcare workers within a single work unit. This study evaluates the risk of burnout in all medical personnel in one perioperative unit.

Methods: We developed an online survey that included demographics, a modified version of the Maslach Burnout Inventory–Human Services Survey, and the Social Support and Personal Coping Survey. Survey constructs (*e.g.*, depersonalization and health) and a global score were calculated. Larger construct and global values were associated with higher risk of burnout. These were separately regressed on role, age, and sex. The global score was then regressed on each of the survey constructs.

Results: Of the 145 responses, 46.2% were physicians (22.8% residents), 43.4% were nurses or nurse anesthetists, and 10.3% were other personnel. After adjusting for sex and age, residents scored higher than other physicians on the following (expected change [95% confidence interval]): global score (1.12 [0.43–1.82]), emotional exhaustion (1.54

What We Already Know about This Topic

- Although burnout is a significant concern among health care providers, a cross-sectional study of burnout among all members of a perioperative unit has not been performed

What This Article Tells Us That Is New

- Among 145 survey respondents using a modified version of the Maslach Burnout Inventory, global burnout scores were higher in physicians than nurse anesthetists and other personnel and highest among residents

[0.44–2.60]), and depersonalization (1.09 [0.23–1.95]). Compared with nonphysicians, residents were 1 U or more higher on these items ($P < 0.05$ in all cases). Residents had higher health (1.49 [0.48–2.50]) and workload (1.23 [0.07–2.40]) values compared with physicians. Better health, personal support, and work satisfaction scores were related to decreased global scores ($P < 0.05$).

Conclusions: Physicians (particularly residents) had the largest global burnout scores, implying increased risk of burnout. Improving overall health, increasing personal support, and improving work satisfaction may decrease burnout among perioperative team members.

* Associate Professor of Clinical Anesthesiology, † Research Coordinator, Department of Anesthesiology, ‡ Professor of Anesthesiology, § Assistant Professor of Biostatistics and Anesthesiology, || Biostatistician III, # Professor of Anesthesiology, Biomedical Informatics, and Medical Education, and Staff Physician, Veterans Affairs Medical Center-Nashville, Tennessee Valley Veterans Affairs Healthcare System, Nashville, Tennessee.

Received from the Departments of Anesthesiology and Biostatistics, Vanderbilt University School of Medicine, Nashville, Tennessee. Submitted for publication February 15, 2010. Accepted for publication June 14, 2010. Support was provided solely from institutional and/or departmental sources. Presented in part at the American Society of Anesthesiologists annual meeting, October 18–22, 2008, Orlando, Florida.

Address correspondence to Dr. Hyman: Department of Anesthesiology, Vanderbilt University Medical Center, 1211 21st Ave S, Medical Arts Building 732, Nashville, Tennessee 37205. steve.hyman@vanderbilt.edu. This article may be accessed for personal use at no charge through the Journal Web site, www.anesthesiology.org.

Copyright © 2010, the American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins. Anesthesiology 2011; 114: 194–204

BURNOUT is a term coined in the early 1970s by Herbert J. Freudenberger, Ph.D., a psychologist and psychoanalyst.^{1,2} The term was given measurable attributes (*i.e.*, emotional exhaustion, depersonalization, and personal accomplishment) by Maslach *et al.*³ in the 1990s. Burnout can lead to poor-quality work, job turnover, and personal and family difficulties. Perioperative clinicians may be at particular risk for burnout given increasing production pressure and staff shortages,³ care of extremely ill patients, and work with extreme responsibility.⁴

For decades, the business community has realized that on-

- ◆ This article is accompanied by an Editorial View. Please see: Shanafelt T: Burnout in anesthesiology: A call to action. *Anesthesiology* 2011; 114:1–2.

the-job factors can affect job performance and satisfaction.⁵⁻⁹ Early studies of physicians did not directly assess burnout; instead, they ascertained career satisfaction or career longevity. For example, Hall *et al.*¹⁰ surveyed emergency department physicians and found the “ten-year survival rate” was 84.9%, (*i.e.*, 15.1% attrition after 10 yr). In 1995, Fields *et al.*¹¹ studied pediatric critical care physicians using the Pines and Aronson burnout scale and found that 14% of respondents were burned out and another 36% were at risk of burnout. Since then, the Maslach Burnout Inventory–Human Services Survey (MBI–HSS) has become the definitive standard for evaluating the risk of job burnout in persons working in human services and health care.³ The Pines and Aronson and the Maslach surveys are the most common burnout measures and are similar in scope (*e.g.*, tiredness and mood) and number (21 and 22, respectively) of their questions. Unlike the Pines and Aronson survey, in which subjects respond to short phrases, the MBI–HSS is written in complete sentences that are specifically geared for subjects who work in human service professions. More recent studies have used the MBI–HSS to evaluate burnout risk in, among others, surgeons,¹² otolaryngologists,¹³ and academic reproductive medicine chairs.¹⁴

These later studies suggested that the risk of burnout and job dissatisfaction has increased in the modern healthcare workforce. Although burnout can occur regardless of work environment,⁴ the risk may be increased by current healthcare work issues, such as financial pressures, reimbursement, and malpractice risk.¹⁴⁻¹⁶ These issues, compounded with inherent personality risk factors,³ can produce a cascade of events leading to significant worker morbidity. These personality traits may include an avoidant coping style and an external locus of control (*i.e.*, a feeling that luck or someone else [rather than one’s self] is responsible for what happens to

an individual). Burnout within a workplace is contagious (*i.e.*, it can spread to coworkers).^{4,13}

Safe surgical care requires high performance by a team of well-trained surgeons, anesthesiologists, nurses, and other personnel; clinical performance can be further mediated by individual factors, including skill, preparation, stress, mood, state of health, and motivation.¹⁷ Performance-shaping factors, including workload, role conflict, lack of community, and value conflicts can predispose to exhaustion, depersonalization, and “inefficacy.”³ Such findings are often signs of burnout and are not unique to anesthesia; however, they can occur in all physician specialties and in nurses and pharmacists. Burnout has been associated with substance abuse,^{16,18,19} mental health issues,²⁰ early retirement,^{21,22} physical illness,^{23,24} and *presenteeism* (a syndrome in which workers are physically present but have low motivation and work effort).²⁵ Presenteeism can cyclically lead to poor job performance and further burnout. In addition to the risk of burnout spreading to other workers, some workers may refuse to work with a burned-out individual.⁴

Most previous studies of burnout in healthcare providers have examined individual specialties; none concurrently assessed burnout in multiple disciplines of the same healthcare team in the same work environment, and none have evaluated the operating room team. We designed the current study to quantify job burnout among perioperative providers in one operating room suite, to identify risk factors associated with job burnout, and to delineate factors that might improve job satisfaction and/or aid in coping with job stress.

Materials and Methods

A survey of 77 questions was developed to assess the magnitude of emotional, mental, and physical exhaustion among

Table 1. Modified MBI–HSS Questions

Construct	MBI–HSS Wording	
	Original	Modified
Depersonalization	I feel I treat some recipients as if they were impersonal objects. I do not really care what happens to some recipients. I feel recipients blame me for some of their problems.	I feel I treat some patients as if they were impersonal objects. I do not really care what happens to some patients. Patients blame me for some of their problems.
Emotional exhaustion	I feel fatigued when I get up in the morning and have to face another day on the job.	I still feel tired when I wake up on workday mornings.
Personal accomplishment	I can easily understand how my recipients feel about things. I deal effectively with the problems of my recipients. I can easily create a relaxed atmosphere with my recipients. I feel exhilarated after working closely with my recipients.	I can easily understand how my patients feel about things. I deal effectively with my patient’s problems. I can easily create a relaxed atmosphere for my patients. I feel exhilarated after working closely with my patients.

HSS = Human Services Survey; MBI = Maslach Burnout Inventory.




Fig. 1. Scores for all social support and personal coping questions. The distributions, as summarized by the 10th through 90th quantiles, of social support and personal coping items and overall constructs are shown. All survey questions are shown as used in the original survey, although those with an asterisk had their tone changed (*i.e.*, from positive to negative) for data analysis.

employees in a busy adult perioperative area. Twenty-two of the questions were the MBI-HSS, whereas the remainder were developed to describe the availability and impact of individuals' social support system and coping strategies.

The Institutional Review Board Behavioral Sciences Committee at Vanderbilt University, Nashville, Tennessee, approved the study. To minimize potential bias, potential participants were not informed of the specific purpose of the study. The act of completing and submitting the question-

naire implied consent. Given the intimate nature of some questions and the potential revelation of undesirable behaviors, it was important that participants not be identifiable. All data were kept completely anonymous to avoid any perceived retribution from participants' superiors. Each questionnaire was numerically coded (to permit follow-up of nonrespondents), but this code was unavailable to the test administrators or data analysts. Consequently, in addition to the general role within the perioperative care environment (*e.g.*, surgeon,

Table 2. Descriptive Summaries by Role

Variable	Role					Overall	P Value
	Resident*	Physician	CRNA	Nurse	Other†		
Participants, No. (%)	34 (23.4)	33 (22.8)	29 (20.0)	34 (23.4)	15 (10.3)	145	NA
Age, y	33.0 [6.0]	51.0 [9.0]	44.0 [9.0]	51.0 [7.0]	44.0 [15.5]	45.0 [16.0]	NA
Male sex, No. (%)	20 (58.8)	23 (69.7)	15 (51.7)	2 (5.9)	3 (20.0)	63 (43.4)	NA
MBI–HSS constructs							
Emotional exhaustion	4.5 [2.6]	3.45 [2.9]	2.45 [2.4]	3.3 [2.1]	3.0 [2.6]	3.4 [2.7]	0.001‡
Depersonalization	3.35 [3.0]	1.45 [1.6]	1.25 [1.8]	1.0 [2.0]	0.8 [1.2]	1.4 [2.0]	<0.001‡
Lack of personal accomplishment	3.75 [1.3]	2.25 [1.2]	3.15 [1.9]	2.1 [1.6]	2.8 [2.4]	2.8 [1.8]	<0.001‡
Global score	4.05 [1.3]	2.45 [1.8]	2.45 [1.4]	2.2 [1.3]	2.1 [1.0]	2.7 [1.8]	<0.001‡
Social support and personal coping constructs							
Health	3.75 [2.4]	2.75 [1.8]	2.55 [3.1]	2.8 [2.6]	2.7 [2.3]	2.9 [2.5]	0.005‡
Personal support	4.55 [1.9]	4.55 [3.0]	3.55 [2.0]	3.0 [3.4]	2.0 [2.8]	4.0 [2.5]	<0.001‡
Professional support	4.75 [2.3]	4.85 [3.5]	3.55 [3.2]	4.8 [2.2]	4.7 [3.0]	4.5 [2.8]	0.698
Workload	5.75 [2.2]	5.05 [2.0]	4.75 [2.0]	4.7 [2.9]	4.7 [2.3]	5.0 [2.3]	0.079
Work satisfaction	4.35 [2.7]	2.85 [2.8]	2.75 [1.2]	3.7 [1.7]	3.5 [1.4]	3.3 [2.4]	0.008‡
Outside activities	2.55 [4.0]	3.05 [3.0]	2.05 [3.0]	1.0 [1.8]	2.0 [2.5]	2.0 [3.0]	0.029‡

Data are given as median [interquartile difference] unless otherwise indicated.

* Physicians with five or fewer years of post–medical school experience were classified as residents for this study. † Clinical job categories in this group included clinical care partner, radiology technician, and one doctoral staff member. ‡ Statistically significant difference between roles by Kruskal–Wallis test. P values were not adjusted for multiple comparisons.

CRNA = certified registered nurse anesthetist; HSS = Human Services Survey; MBI = Maslach Burnout Inventory; NA = not applicable.

nurse, or anesthesia provider), we did not collect any data (e.g., resident status) that could potentially be used to identify individual respondents. Participants accessed a Web site linked to a secure database (Vovici, Dulles, VA).

Study Population

Participants were employees with patient contact responsibilities in an adult perioperative suite (including preoperative, intraoperative, and postoperative areas). Eligible participants included nurses, certified registered nurse anesthetists, surgeons (resident and attending), anesthesiologists (resident and attending), scrub technicians, and other nondegreed clinical care providers. Medical students and student nurse anesthetists were excluded. In this study, *residents* were defined as physicians with five or fewer years of post–medical school experience.

We specifically targeted the unit's primary healthcare professional workforce, 250 potential participants, by e-mail and asked them to complete the Web-based survey. To maximize response rate, we sent three sequential e-mails to the participants. In addition, we placed recruitment posters in strategic locations (i.e., break rooms, locker rooms, control rooms, and nurses' stations).

Survey Questions

Modified Maslach Constructs. The MBI–HSS evaluates the three aspects of burnout (i.e., emotional exhaustion, depersonalization, and personal accomplishment). We altered the original MBI–HSS in three ways:

1. The possible responses for the original Maslach survey ranged from never (0) to every day (6) and asked the respondent to recall information from the past year. We focused on the past 2–4 weeks and ranked them on a scale of lowest (0) to highest (9) to provide a finer resolution of the responses.
2. We slightly altered the wording of eight items from the original MBI–HSS instrument to make it more applicable to perioperative care providers (table 1).
3. The MBI–HSS items (or their modified counterparts) were intertwined with other questions and were not in their original order.

To assess the effect of these modifications on the reliability of each construct, Cronbach's α statistics were calculated on the collected survey data. The α values for emotional exhaustion, depersonalization, and personal accomplishment were 0.88, 0.77, and 0.82, respectively, which indicates that even with the modifications the questions measured similar structures. Regardless, these modifications negated the applicability of the original instrument's scoring metrics. The average of the three constructs was used to define a "burnout global score." Because the tone of emotional exhaustion and depersonalization is negative, the personal accomplishment construct was transformed into a lack of personal accomplishment construct by subtracting nine from the personal accomplishment score because nine was the maximum item score in our version of the MBI–HSS.

Social Support and Personal Coping Constructs. The remainder of the survey was composed of questions intended to delineate individual's coping strategies and social support sys-

Table 3. Multiple Linear Regression Models for MBI–HSS and Social Support and Personal Coping Constructs

Outcomes	Resident (Intercept)	Physician	CRNA
MBI–HSS constructs			
Global score	3.94 (3.49, 4.39)	–1.12 (–1.82, –0.43)*	–1.26 (–1.88, –0.63)*
Emotional exhaustion	4.79 (4.07, 5.51)	–1.54 (–2.64, –0.44)*	–1.86 (–2.86, –0.87)*
Depersonalization	3.24 (2.68, 3.81)	–1.09 (–1.95, –0.23)*	–1.39 (–2.17, –0.62)*
Low personal accomplishment	3.79 (3.23, 4.34)	–0.74 (–1.59, 0.11)	–0.51 (–1.28, 0.25)
Social support and personal coping constructs			
Health	4.13 (3.47, 4.79)	–1.49 (–2.50, –0.48)*	–1.37 (–2.28, –0.47)*
Outside activities	3.19 (2.17, 4.20)	0.92 (–0.63, 2.47)	0.00 (–1.39, 1.40)
Personal support	4.45 (3.76, 5.13)	–0.40 (–1.45, 0.65)	–1.19 (–2.14, –0.25)*
Professional support	5.39 (4.56, 6.23)	0.36 (–0.92, 1.64)	–0.32 (–1.48, 0.83)
Work satisfaction	4.61 (3.94, 5.29)	–0.98 (–2.01, 0.05)	–1.29 (–2.22, –0.36)*
Workload	5.76 (5.00, 6.53)	–1.23 (–2.40, –0.07)*	–0.98 (–2.03, 0.07)

Data are given as parameter estimates (95% confidence intervals).

* Values that were statistically significant at the 0.05 level.

CRNA = certified registered nurse anesthetist; HSS = Human Services Survey; MBI = Maslach Burnout Inventory.

tem. These questions occurred in the following six groupings: health ($n = 15$), personal ($n = 2$), professional support ($n = 6$), workload ($n = 3$), work satisfaction ($n = 10$), and outside activities ($n = 1$). Each was defined as the average score of the construct-specific items after adjusting for tone. Most of the questions were originally phrased in a negative tone; thus, any item with a positive tone was adjusted to a negative tone for the analysis so that items could ultimately be grouped into a single score. For example, “My job satisfies me economically,” with a score of 4, was replaced with “My job does not satisfy me economically,” with a score of 5. A complete listing of survey questions with the original wording is in the figure.

Statistical Analysis

Descriptive summaries were calculated for age, sex, MBI–HSS constructs, and global scores. Counts and percentages were used for categorical variables, and medians and interquartile differences (75th–25th) were used for continuous variables. To inspect how burnout scores and the social support and personal coping constructs differed among the various roles, the Kruskal–Wallis test was applied.

We examined the association between role, sex, and age and the MBI–HSS and social support and personal coping scores using a series of linear regression models. Role and sex were coded as indicator variables, with female residents being defined as the reference group. Although any group could have been used as the reference group in the models, based on previous studies, younger physicians might be expected to have higher burnout scores; thus, it seemed logical to use them as the reference. Age was modeled as a linear term and was centered at 30 yr, roughly the median age of the resident sample. The resultant age estimate and its SE were scaled to reflect 10-yr increments. Modeling assumptions were evaluated by visual inspection of residual and QQ plots. All data analyses were performed using computer software (R Version 2.9.0; R Development Core Team, Vienna, Austria).²⁶

Results

During the survey period (February–July 2007), 145 surveys were completed. If only the 250 surgeons, nurses, and anesthesia providers who work in this area regularly and were targeted for survey participation were the potential respondents, then the response rate was 58%. However, approximately 150 other personnel (mostly anesthesia providers) occasionally work in this operating room suite and could have seen the “participation encouragement” posters. If all of these “occasional” personnel saw the poster and declined to participate, the response rate was closer to 36%.

Of survey respondents, physicians (attending and resident) and nurses (registered nurses, licensed practical nurses, or certified registered nurse anesthetists) constituted 46% and 43% of the responses, respectively. Female respondents composed 57% of the sample. Most physicians were men (64% of 67), and most nurses were women (73% of 63). The age (median [interquartile difference]) across all roles was 45 [16] yr. These demographic summaries, stratified by role, are presented in table 2, along with summaries each MBI–HSS construct and the global score. Median scores for emotional exhaustion, depersonalization, and low personal accomplishment were as follows: 3.4, 2.7, 1.4, 2.0, and 2.8, 1.8, respectively. These scores and the global score differed significantly among the roles ($P < 0.001$). The median scores for social support and personal coping items were as follows: health (2.9, 2.5), personal support (4.0, 2.5), professional support (4.5, 2.8), workload (5.0, 2.3), work satisfaction (3.3, 2.4), and outside activities (2.0, 3.0). Significant differences by role were found for all items ($P < 0.05$), except professional support and workload.

Social support and personal coping summaries are presented in the figure, along with the item-specific responses for each category (e.g., health and personal support). The highest median scores (all >5 of 9) were observed in the following subitems: feeling worn out at the end of a shift,

Table 3. Continued

Nurse	Other	Male Sex	Aged 10 yr
-1.24 (-1.95, -0.52)*	-1.26 (-2.01, -0.51)*	0.04 (-0.38, 0.46)	-0.21 (-0.48, 0.05)
-1.87 (-3.02, -0.73)*	-2.00 (-3.20, -0.81)*	-0.23 (-0.90, 0.44)	0.12 (-0.30, 0.55)
-1.18 (-2.08, -0.29)*	-1.95 (-2.89, -1.02)*	0.33 (-0.20, 0.85)	-0.41 (-0.74, -0.08)*
-0.65 (-1.53, 0.23)	0.18 (-0.74, 1.10)	0.01 (-0.50, 0.53)	-0.36 (-0.68, -0.03)*
-1.53 (-2.58, -0.49)*	-1.74 (-2.84, -0.65)*	-0.36 (-0.97, 0.25)	-0.11 (-0.28, 0.49)
-0.33 (-1.94, 1.28)	0.19 (-1.49, 1.86)	0.77 (-0.17, 1.71)	-0.57 (-1.16, 0.02)
-1.48 (-2.56, -0.39)*	-1.97 (-3.10, -0.83)*	0.85 (0.21, 1.48)*	-0.18 (-0.58, 0.22)
-0.13 (-1.46, 1.20)	0.10 (-1.29, 1.48)	-0.74 (-1.52, 0.04)	-0.39 (-0.88, 0.10)
-0.66 (-1.73, 0.42)	-0.74 (-1.86, 0.37)	-0.49 (-1.11, 0.14)	-0.11 (-0.51, 0.28)
-1.23 (-2.44, -0.02)*	-1.34 (-2.61, -0.08)*	-0.18 (-0.89, 0.53)	-0.11 (-0.34, 0.55)

insufficient departmental emotional support, and not having control over one's workload.

Table 3 presents parameter estimates and 95% confidence intervals for the following two linear regression models: (1) MBI-HSS constructs on role, age, and sex; and (2) social support and personal coping constructs on role, age, and sex. Based on the coding of role, age, and sex, the resident column (the intercept of each model) corresponds to female residents aged 30 yr. Other values in this table correspond to expected changes in MBI-HSS or social support and personal coping scores associated with (1) another role *versus* the resident role, (2) being male *versus* female, and (3) 10-yr age changes. Residents had consistently higher MBI-HSS scores than any other clinical role. They scored significantly higher than other physicians on global score, emotional exhaustion, and depersonalization constructs but not on low personal accomplishment. Compared with the nonphysician roles, resident scores were often greater than 1 U higher on the global score, emotional exhaustion, and depersonalization ($P < 0.05$ in all cases) but not low personal accomplishment. Although sex was not independently predictive of MBI-HSS construct scores, age had a clear impact on depersonalization and low personal accomplishment. The estimated change per 10-yr increase in age was -0.41 (-0.74 to -0.08) and -0.36 (-0.68 to -0.03) for depersonalization and low personal accomplishment, respectively. Age was not a strong predictor of either the global score or emotional exhaustion.

Residents and other physicians scored similarly on social support and personal coping items other than health and workload. Regression parameter estimates associated with being a physician (*versus* being a resident) were -1.49 (-2.50 to -0.48) for the health item and -1.23 (-2.40 to -0.07) for the workload item. Health and personal support scores for residents were at least 1 U higher than those for other nonphysician roles ($P < 0.05$), and results were nearly as strong for the workload item. Furthermore, residents

scored 1.29 (0.36–2.22) U higher than nurse anesthetists on work satisfaction.

When the global score was regressed on the social support and personal coping items, the health, personal support, and work satisfaction items contributed to a higher global score ($P < 0.05$). Health had the strongest impact, with a 1-U increase in the health score corresponding to a 0.32 (0.21–0.43) U increase in global score. There were smaller, but still significant, effects of personal support and work satisfaction on the global score.

Discussion

In this study, we used a questionnaire consisting of a modified MBI-HSS and our own questions (social support and personal coping) to evaluate the risk of burnout among healthcare providers in a single surgical unit. Physicians (particularly residents) seem to be at greatest risk. Age was associated with the depersonalization and low personal accomplishment items but sex was not. After adjusting for role, age was not associated with any of the social support and personal coping items, but sex was related to personal support. Health, followed by work satisfaction and personal support, had the greatest effect on global score. This implies that maintaining good health, along with the availability of family and friends, contributes to more favorable personal coping strategies.

Maslach's definition of burnout is an increased level of emotional exhaustion and depersonalization, with low levels of personal accomplishment.³ Because burnout "is an individual experience that is specific to the work context," six factors are believed to be "prime correlates" for the development of burnout: job, occupational, organizational, demographic, and personality characteristics and job attitudes. At-risk negative job characteristics may involve workload (overwork or boredom), conflict, diminished resources or social support, and lack of input or feedback. The "classical"

Table 4. Comparison of Studies of Burnout in Physicians

Study	Specialty Studied	Country of Study Population	Instrument(s) Used
Hall <i>et al.</i> , ¹⁰ 1992	Emergency medicine	United States	P&A
Lloyd <i>et al.</i> , ⁴⁰ 1994	Emergency medicine	Canada	MBI, RDAS, EPJS, or T.O.
Fields <i>et al.</i> , ¹¹ 1995	Pediatric critical care	United States	P&A or T.O.
Guntupalli and Fromm, ³⁶ 1996	Intensive care	United States	MBI or T.O.
Ramirez <i>et al.</i> , ¹⁵ 1996	Gastroenterology	United Kingdom	MBI, T.O., or GHQ-12
	Surgery	United Kingdom	MBI, T.O., or GHQ-12
	Radiology	United Kingdom	MBI, T.O., or GHQ-12
	Oncology	United Kingdom	MBI, T.O., or GHQ-12
Campbell <i>et al.</i> , ¹² 2001	Surgery (multispecialty)	United States	MBI
Hoff <i>et al.</i> , ³³ 2001	Hospitalist	United States	P&A
Gabbe <i>et al.</i> , ¹⁴ 2002	Reproductive medicine (chairs)	United States	MBI,* APS, APPS, or SSS
Sargent <i>et al.</i> , ³⁸ 2004	Orthopedics	United States	MBI, RDAS, or GHQ-12
Nyssen <i>et al.</i> , ³¹ 2003; and Nyssen and Hansez, ⁶⁰ 2008	Anesthesia	Belgium	T.O., PSSM-A, WOCCQ, or SRPHS
Bertges Yost <i>et al.</i> , ³⁴ 2005	Transplantation	United States	MBI, T.O., or SCI
Johns and Ossoff, ¹³ 2005; and Golub <i>et al.</i> , ⁵⁰ 2007	Otolaryngology	United States	MBI,* APS, APPS, or SSS
Garelick <i>et al.</i> , ²⁴ 2007	All specialties	United Kingdom	MBI, Core-OM, CORE-A, CORE- workplace, BSI, or GSI
Raggio and Malacarne, ³⁷ 2007	Critical care	Italy	MBI, PMS, or DMI
Magnavita <i>et al.</i> , ⁴⁵ 2008	Radiology	Italy	MBI, T.O., GHQ-12, KJCS, SERI, WJS, or GADS
Sharma <i>et al.</i> , ⁶¹ 2008	Colorectal	United Kingdom	MBI, T.O., GHQ-12, or TCQ
Embriaco <i>et al.</i> , ³⁵ 2007	Critical care	Italy	MBI, T.O., SAPSII, or CESDS
Current study	Perioperative providers	United States	MBI-HSS

* Used a shortened MBI with a reliability test.

APS = Assessment of Professional Stressors; APPS = Assessment of Personal and Professional Satisfaction; BSI = Brief System Inventory; CESDS = Centers for Epidemiological Studies Depression Scale; CORE = Clinical Outcomes in Routine Evaluation; DMI = Defense Mechanism Inventory; EPJS = Emergency Physicians Job Satisfaction Instrument; GADS = Goldberg's Anxiety and Depression Scales; GHQ-12 = General Health Questionnaire-12; GSI = Global Severity Index; KJCS = Karasek's Job Content Questionnaire; MBI = Maslach Burnout Index; NA = not applicable; NR = not reported; OM = Outcome Measure; P&A = Pines and Aronson; PMS = Profile of Mood States; PSSM-A = Psychological State of Stress Measures; RDAS = Revised Dyadic Adjustment Scale; SAPSII = Simplified Acute Physiology Scale; SCI = Surgeon Coping Inventory; SERI = Siegert's Effort-Reward Imbalance; SRPHS = Self-reported Physical Health Scale; SSS = Spousal Support Survey; TCQ = The Coping Questionnaire; T.O. = Their Own Questionnaire; WJS = Warr's Job Satisfaction; WOCCQ = Working Conditions and Control Questionnaire.

at-risk organizational profile is one in which there is a steep hierarchy and more is demanded of employees and less is given by the employer, such as that seen with downsizing or mergers. At-risk demographics include younger adults with more education and busy unmarried people (especially those who have never been married). At-risk personality characteristics are low hardiness (*i.e.*, without "involve[ment] in daily activities, a sense of control over events, and openness to change," poor self-esteem, and external locus of control [the "victim" mentality]). An at-risk job attitude is when people have unrealistically high expectations for their job. It is prudent to identify those who are at risk for burnout because it can

contribute to worsening job performance (*i.e.*, absenteeism, job turnover, decreased productivity, and negative effect on coworkers) and health issues (*i.e.*, substance abuse and mental and physical problems).^{4,20,23,27-29}

Interpretation of the literature regarding burnout is challenging (table 4). Some articles^{11,30-32} use the term burnout and assert its presence in their participants but are unclear about their measurement methods or criteria. Although many older studies used the Pines and Aronson instrument to assess burnout,^{10,33} most current studies use the MBI-HSS as their primary measurement tool.^{12-15,22,34-38} Our study used a variant of the MBI-HSS tailored to our periop-

Table 4. Continued

Attending				Resident				
Moderate to High Emotional Exhaustion, % (% High)	Moderate to High Depersonalization, % (% High)	Moderate to Low Personal Accomplishment, % (% Low)	Medium to High Burnout, % (% High)	Moderate to High Emotional Exhaustion, % (% High)	Moderate to High Depersonalization, % (% High)	Moderate to Low Personal Accomplishment, % (% Low)	Moderate to High Burnout, % (% High)	Chair: Moderate to High Burnout % (% High)
NA	NA	NA	(23)	NA	NA	NA	NA	NA
46 (29)	93 (61)	79 (16)	NA	NA	NA	NA	NA	NA
NA	NA	NA	50	NA	NA	NA	NA	NA
(29)	(20)	(59)	NA	NA	NA	NA	NA	NA
(31)	(28)	(38)	NR	NA	NA	NA	NA	NA
(27)	(19)	(38)	NR	NA	NA	NA	NA	NA
(33)	(21)	(49)	NR	NA	NA	NA	NA	NA
(35)	(27)	(37)	NR	NA	NA	NA	NA	NA
63 (32)	37 (13)	17 (4)	NA	NA	NA	NA	NA	NA
NA	NA	NA	37 (13)	NA	NA	NA	NA	NA
80 (56)	65 (36)	39 (21)	NA	NA	NA	NA	NA	92 (4)
Low	Low	High	Not high	High	High	Low	High	NA
NA	NA	NA	88 (22)	NA	NA	NA	NA	NA
61 (38)	49 (27)	40 (16)	NR	NA	NA	NA	NA	NA
35	20	68	Chair	35 (26)	20 (13)	68 (47)	86 (10)	84 (3)
(66)	(39)	(40)	18	NA	NA	NA	NA	NA
(36)	(56)	(28)	NR	NA	NA	NA	NA	NA
NA	NA	NA	24–38	NA	NA	NA	NA	NA
(31)	(17)	(27)	NA	NA	NA	NA	NA	NA
NR	NR	NR	46 (30)	NA	NA	NA	NA	NA
NA	NA	NA	NA	NA	NA	NA	NA	NA

erative participant population. Although most of the studies in table 4 used Maslach or Pines and Aronson, some did not. Several articles used the General Health Questionnaire-12 as an additional instrument, but most either created their own questions or used a unique instrument. Future research on clinician burnout should strive for consistency in study design and choice of study instruments. We recommend the MBI-HSS.

In agreement with some studies, we were unable to detect differences in the MBI-HSS items between men and women,^{22,33,37} but men were more likely to have a higher personal support score (table 3) after adjusting for age

and role. Other researchers^{10,13,15,30–32,36,38–40} have not commented on this correlation, but Fields *et al.*¹¹ and Guntupalli and Fromm³⁶ have reported increased burnout in female intensive care physicians. Gabbe *et al.*¹⁴ identified increased emotional exhaustion and decreased personal accomplishment (both bad) and decreased depersonalization (good) in female reproductive medicine chairs but did not identify them as burned out. If female healthcare professionals have more burnout characteristics than male counterparts but also have less support, they are at greater risk and interventions may need to be targeted to prevent them from burning out.

Overall, our physicians appeared to be at higher risk for burnout than were nurses. This finding is consistent with two studies of nurses^{41,42} and two studies that included nurses.^{22,37} Other nurse-centered studies^{43,44} have reported a high burnout rate, but their data must be viewed skeptically because of how they determined burnout (as discussed later). Some physician specialties appear to have a relatively low risk of burnout,^{12,22} but others may have an enormous problem.^{15,31,40,45} Surprisingly, specialties that might be considered high stress because of the nature of the work (surgery, transplantation, or orthopedics) did not report excessively high burnout scores,^{12,14,34,38} whereas intensivists and anesthesiologists had a greater incidence.^{31,35–37} It seems counterintuitive, but other specialties for which daily work might not at first seem highly stressful (*e.g.*, otolaryngology and radiology) have been at higher risk for burnout.^{13,15,45}

Some researchers, based on studies of physicians^{13–15,34,36} and nurses,⁴¹ suggest that personal accomplishment is a buffering element. Even faced with overwhelming stress and job dissatisfaction, burnout risk decreases if individuals feel like their professional activities are for a good purpose. Aiken *et al.*⁴³ and Erickson and Grove⁴⁴ reported a burnout rate in nurses of 38–42%, but both studies equated burnout with high levels of emotional exhaustion. These studies failed to consider the potential protective effects of personal accomplishment in their analysis. Despite more than 43,000 subjects, their actual rate of burnout could be lower (if personal accomplishment were high) or higher (if personal accomplishment were low). Future studies will be needed to understand whether and how personal accomplishment really protects high-risk nurses and physicians (*i.e.*, high emotional exhaustion and depersonalization) from manifesting burnout.

Given the prevalence of burnout in health care, more investigators could be working on mitigation strategies. Scandinavian researchers^{46,47} have successfully used peer support groups to treat burned-out providers (*i.e.*, physicians, nurses, therapists, and social workers), but only a few U.S. articles have proactively dealt with burnout. Elpern and Silver⁴⁸ reviewed burnout in the intensive care unit and provided guidelines for an organization to manage burnout. Dunn *et al.*⁴⁹ describes a proactive data-guided program (using validated instruments) that a private primary care group adopted to “enhance physician and organizational well-being.”

Perhaps the most compelling aspect of our results is in our resident category (*i.e.*, the respondents who had the least experience and were the lowest in the organizational physician hierarchy). These physicians reported more negative characteristics than did any other clinicians. Although physicians in general appeared to be worse off relative to nurses in the areas of personal and professional support and outside activities (tables 2 and 3), residents were most affected in the areas of health, personal support, and workload. Residents scored more poorly in the global score,

emotional exhaustion, and depersonalization but were notably worse off in personal accomplishment ($P < 0.05$, table 3). Our findings are similar to other studies with residents^{38,50} and younger physicians, a pattern also seen in younger nurses.^{41,43,44}

The general demographic of the burned-out employee matches that of medical trainees (young, single, well educated, and less in control of work life).³ Dyrbye *et al.* reported that nearly half of medical students at major schools in the United States are burned out⁵¹ and that negative personal life events affected their burnout rate.⁵² Burnout remains a significant problem in residents^{50,53–57} and may affect most of that group. Early intervention in these two groups alone might yield happier physicians who lead longer and more productive careers. Unfortunately, the literature on effective interventions is sparse. As with practicing physicians, few studies adequately address the issue of medical trainee burnout and its diagnosis, treatment, or prevention. McCray *et al.*,⁵⁸ reviewing literature from 1996 to 2007, found 190 articles and analyzed 129 articles. Of these studies, only 9 pertained specifically to therapy for residents or medical students and only 2 were randomized controlled trials. Sadly, none met the Strength of Recommendation Taxonomy⁵⁹ criteria for an “A” grade (“recommendation based on consistent and good-quality” evidence). There are no data on the effects of resident work hour restrictions on burnout on either the residents themselves or on the attending physicians or nurse practitioners who have had to pick up their work.

Another important consideration is what effect burnout has on the quality of patient care (*i.e.*, do affected individuals make more medical errors?). Unfortunately, the literature is confusing and inadequate for answering this question. West *et al.*⁵⁷ and Shanafelt *et al.*⁵⁵ report increased medical errors in burned-out residents, but Fahrenkopf *et al.*⁵³ did not. The problem with these studies is their reliance on self-reported errors. A change in the reported incidence of errors could be the result of a change in reporting (*e.g.*, burned-out or depressed physicians might be more or less likely to report errors), a change in the recognition of errors, or a change in the actual occurrence of errors. Thus, future research will be required to ascertain the nature of the relationship between clinician burnout and patient safety.

The term used by Maslach *et al.*³ for the opposite of job burnout is job engagement and involves the proper alignment of the individual’s own characteristics with his or her job characteristics. People can be taught new coping strategies, but workplace issues such as equity and fairness must also be addressed. Examples of positive healthcare workplace interventions in medicine might include flexible work hours, greater interdisciplinary clinical decision making,⁴⁸ and meetings to allow workers to express their concerns.⁴⁹ Although organizational changes are more expensive and more difficult to implement than are those for individuals, mean-

ingful change can only occur if both areas are adequately addressed.

This study has limitations. Although Shanafelt *et al.*⁵⁵ believe that burned-out or depressed physicians are more likely than their unaffected counterparts to report suboptimal care, it is possible that both the most and the least severely affected individuals in our population did not complete our survey (*i.e.*, response bias). It is unknown if smaller or greater differences between roles, sexes, and age groups would be observed if willingness to complete the survey was based solely on a physician's burnout status. Additional studies will be needed to try to capture the widest possible array of individuals at risk.

A second limitation is our modifications to the MBI-HSS items. Based on Cronbach's α statistics (>0.75), these modifications had little impact on the internal reliability of each construct. Nevertheless, the actual score values cannot be directly compared with previously published Maslach benchmarks. Finally, in our well-intended attempt to maintain complete anonymity, we were unable to obtain important data relating to the educational status or specialty (*e.g.*, surgeon or anesthesiologist) of our physician respondents.

In conclusion, we found that, as opposed to the nurses and nurse anesthetists, physicians (particularly those who were younger) had higher levels of depersonalization and emotional exhaustion and are at a higher risk of burnout. Given the known risks associated with job burnout, these data suggest the need for greater attention to this phenomenon in the modern perioperative environment. Maslach *et al.*³ suggest that a successful job-to-person match depends on workload, individual control, appropriate rewards, a sense of community, a sense of fairness, and common values. These items must all be in balance to achieve job engagement, the antithesis of burnout.³ Perhaps individuals can be taught to cope with their job demands and, thus, reduce burnout. The impact of recent nationwide efforts to reengineer health care, incentivize out poor quality, and reduce costs must consider burnout risks in all medical and allied health personnel. Future studies need to capture more data in other surgical and nonsurgical units to determine whether our results are generalizable, to identify the highest-risk groups and contributory factors, and to evaluate prevention and treatment interventions.

References

- Freudenberger HJ: The staff burn-out syndrome in alternative institutions. *Psychotherapy: Theory, Research, and Practice*. 1975; 12(1):73-82
- Freudenberger HJ: Burn-out: Occupational hazard of the child care worker. *Child Care Q* 1977; 6:90-9
- Maslach C, Schaufeli WB, Leiter MP: Job burnout. *Annu Rev Psychol* 2001; 52:397-422
- Felton JS: Burnout as a clinical entity: Its importance in health care workers. *Occup Med (Lond)* 1998; 48:237-50
- Raymond CA: Mental stress: "Occupational injury" of 80s that even pilots can't rise above. *JAMA* 1988; 259:3097-8
- Little LF, Gaffney IC, Rosen KH, Bender MM: Corporate instability is related to airline pilots' stress symptoms. *Aviat Space Environ Med* 1990; 61:977-82
- Girodo M: The psychological health and stress of pilots in a labor dispute. *Aviat Space Environ Med* 1988; 59:505-10
- Dell'Erba G, Venturi P, Rizzo F, Porcu S, Pancheri P: Burnout and health status in Italian air traffic controllers. *Aviat Space Environ Med* 1994; 65:315-22
- Singh RG: Relationship between occupational stress and social support in flight nurses. *Aviat Space Environ Med* 1990; 61:349-52
- Hall KN, Wakeman MA, Levy RC, Khoury J: Factors associated with career longevity in residency-trained emergency physicians. *Ann Emerg Med* 1992; 21:291-7
- Fields AI, Cuedon TT, Brasseux CO, Getson PR, Thompson AE, Orłowski JP, Youngner SJ: Physician burnout in pediatric critical care medicine. *Crit Care Med* 1995; 23:1425-9
- Campbell DA Jr, Sonnad SS, Eckhauser FE, Campbell KK, Greenfield LJ: Burnout among American surgeons. *Surgery* 2001; 130:696-702; discussion, 702-5
- Johns MM 3rd, Ossoff RH: Burnout in academic chairs of otolaryngology: Head and neck surgery. *Laryngoscope* 2005; 115:2056-61
- Gabbe SG, Melville J, Mandel L, Walker E: Burnout in chairs of obstetrics and gynecology: Diagnosis, treatment, and prevention. *Am J Obstet Gynecol* 2002; 186:601-12
- Ramirez AJ, Graham J, Richards MA, Cull A, Gregory WM: Mental health of hospital consultants: The effects of stress and satisfaction at work. *Lancet* 1996; 347:724-8
- Spickard A Jr, Gabbe SG, Christensen JF: Mid-career burnout in generalist and specialist physicians. *JAMA* 2002; 288:1447-50
- Weinger MB, Englund CE: Ergonomic and human factors affecting anesthetic vigilance and monitoring performance in the operating room environment. *ANESTHESIOLOGY* 1990; 73:995-1021
- Brooke D, Edwards G, Taylor C: Addiction as an occupational hazard: 144 Doctors with drug and alcohol problems. *Br J Addict* 1991; 86:1011-6
- Juntunen J, Asp S, Olkinuora M, Aarimaa M, Strid L, Kauttu K: Doctors' drinking habits and consumption of alcohol. *BMJ* 1988; 297:951-4
- Ahola K, Kivimaki M, Honkonen T, Virtanen M, Koskinen S, Vahtera J, Lonnqvist J: Occupational burnout and medically certified sickness absence: A population-based study of Finnish employees. *J Psychosom Res* 2008; 64:185-93
- Gardulf A, Soderstrom IL, Orton ML, Eriksson LE, Arnetz B, Nordstrom G: Why do nurses at a university hospital want to quit their jobs? *J Nurs Manag* 2005; 13:329-37
- Sharma A, Sharp DM, Walker LG, Monson JR: Stress and burnout among colorectal surgeons and colorectal nurse specialists working in the National Health Service. *Colorectal Dis* 2008; 10:397-406
- Borritz M, Rugulies R, Christensen KB, Villadsen E, Kristensen TS: Burnout as a predictor of self-reported sickness absence among human service workers: Prospective findings from three year follow up of the PUMA study. *Occup Environ Med* 2006; 63:98-106
- Garellick AI, Gross SR, Richardson I, von der Tann M, Bland J, Hale R: Which doctors and with what problems contact a specialist service for doctors? A cross sectional investigation. *BMC Med* 2007; 5:26
- Koopman C, Pelletier KR, Murray JF, Sharda CE, Berger ML, Turpin RS, Hackleman P, Gibson P, Holmes DM, Bendel T: Stanford presenteeism scale: Health status and employee productivity. *J Occup Environ Med* 2002; 44:14-20
- R Development Core Team: R: A Language and Environment for Statistical Computing. Vienna, Austria, R Foundation for Statistical Computing, 2009

27. Middaugh DJ: Presenteeism: Sick and tired at work. *Dermatol Nurs* 2007; 19:172-3, 185
28. Pilette PC: Presenteeism in nursing: A clear and present danger to productivity. *J Nurs Adm* 2005; 35:300-3
29. Toppinen-Tanner S, Ojajarvi A, Vaananen A, Kalimo R, Japinen P: Burnout as a predictor of medically certified sick-leave absences and their diagnosed causes. *Behav Med* 2005; 31:18-27
30. Kisa S, Kisa A: Job dissatisfaction among public hospital physicians is a universal problem: Evidence from Turkey. *Health Care Manag (Frederick)* 2006; 25:122-9
31. Nyssen AS, Hansez I, Baelle P, Lamy M, De Keyser V: Occupational stress and burnout in anaesthesia. *Br J Anaesth* 2003; 90:333-7
32. Gaba DM, Howard SK: Patient safety: Fatigue among clinicians and the safety of patients. *N Engl J Med* 2002; 347:1249-55
33. Hoff TH, Whitcomb WF, Williams K, Nelson JR, Cheesman RA: Characteristics and work experiences of hospitalists in the United States. *Arch Intern Med* 2001; 161:851-8
34. Bertges Yost W, Eshelman A, Raoufi M, Abouljoud MS: A national study of burnout among American transplant surgeons. *Transplant Proc* 2005; 37:1399-401
35. Embriaco N, Azoulay E, Barrau K, Kentish N, Pochard F, Loundou A, Papazian L: High level of burnout in intensivists: Prevalence and associated factors. *Am J Respir Crit Care Med* 2007; 175:686-92
36. Guntupalli KK, Fromm RE Jr: Burnout in the internist-intensivist. *Intensive Care Med* 1996; 22:625-30
37. Raggio B, Malacarne P: Burnout in intensive care unit. *Minerva Anesthesiol* 2007; 73:195-200
38. Sargent MC, Sotile W, Sotile MO, Rubash H, Barrack RL: Stress and coping among orthopaedic surgery residents and faculty. *J Bone Joint Surg Am* 2004; 86-A:1579-86
39. Kushnir T, Cohen AH: Job structure and burnout among primary care pediatricians. *Work* 2006; 27:67-74
40. Lloyd S, Streiner D, Shannon S: Burnout, depression, life and job satisfaction among Canadian emergency physicians. *J Emerg Med* 1994; 12:559-65
41. Kilfedder CJ, Power KG, Wells TJ: Burnout in psychiatric nursing. *J Adv Nurs* 2001; 34:383-96
42. Sadovich JM: Work excitement in nursing: An examination of the relationship between work excitement and burnout. *Nurs Econ* 2005; 23:91-6, 55
43. Aiken LH, Clarke SP, Sloane DM, Sochalski JA, Busse R, Clarke H, Giovannetti P, Hunt J, Rafferty AM, Shamian J: Nurses' reports on hospital care in five countries. *Health Aff (Millwood)* 2001; 20:43-53
44. Erickson R, Grove WJC: Why emotions matter: age, agitation, and burnout among registered nurses. *Online J Issues Nurs* 2008; 13
45. Magnavita N, Fileni A, Magnavita G, Mammi F, Mirk P, Roccia K, Bergamaschi A: Work stress in radiologists: A pilot study. *Radiol Med* 2008; 113:329-46
46. Peterson U, Bergstrom G, Samuelsson M, Asberg M, Nygren A: Reflecting peer-support groups in the prevention of stress and burnout: Randomized controlled trial. *J Adv Nurs* 2008; 63:506-16
47. Ro KE, Gude T, Tyssen R, Aasland OG: Counselling for burnout in Norwegian doctors: One year cohort study. *BMJ* 2008; 337:a2004
48. Elpern EH, Silver MR: Improving outcomes: Focus on workplace issues. *Curr Opin Crit Care* 2006; 12:395-8
49. Dunn PM, Arnetz BB, Christensen JF, Homer L: Meeting the imperative to improve physician well-being: Assessment of an innovative program. *J Gen Intern Med* 2007; 22:1544-52
50. Golub JS, Weiss PS, Ramesh AK, Ossoff RH, Johns MM 3rd: Burnout in residents of otolaryngology-head and neck surgery: A national inquiry into the health of residency training. *Acad Med* 2007; 82:596-601
51. Dyrbye LN, Thomas MR, Massie FS, Power DV, Eacker A, Harper W, Durning S, Moutier C, Szydlo DW, Novotny PJ, Sloan JA, Shanafelt TD: Burnout and suicidal ideation among U.S. medical students. *Ann Intern Med* 2008; 149:334-41
52. Dyrbye LN, Thomas MR, Huntington JL, Lawson KL, Novotny PJ, Sloan JA, Shanafelt TD: Personal life events and medical student burnout: A multicenter study. *Acad Med* 2006; 81:374-84
53. Fahrenkopf AM, Sectish TC, Barger LK, Sharek PJ, Lewin D, Chiang VW, Edwards S, Wiedermann BL, Landrigan CP: Rates of medication errors among depressed and burnt out residents: Prospective cohort study. *BMJ* 2008; 336:488-91
54. Sawyer RG, Tribble CG, Newberg DS, Pruett TL, Minasi JS: Intern call schedules and their relationship to sleep, operating room participation, stress, and satisfaction. *Surgery* 1999; 126:337-42
55. Shanafelt TD, Bradley KA, Wipf JE, Back AL: Burnout and self-reported patient care in an internal medicine residency program. *Ann Intern Med* 2002; 136:358-67
56. Tyssen R, Vaglum P, Gronvold NT, Ekeberg O: The impact of job stress and working conditions on mental health problems among junior house officers: A nationwide Norwegian prospective cohort study. *Med Educ* 2000; 34:374-84
57. West CP, Huschka MM, Novotny PJ, Sloan JA, Kolars JC, Habermann TM, Shanafelt TD: Association of perceived medical errors with resident distress and empathy: A prospective longitudinal study. *JAMA* 2006; 296:1071-8
58. McCray LW, Cronholm PF, Bogner HR, Gallo JJ, Neill RA: Resident physician burnout: Is there hope? *Fam Med* 2008; 40:626-32
59. Ebell MH, Siwek J, Weiss BD, Woolf SH, Susman J, Ewigman B, Bowman M: Strength of recommendation taxonomy (SORT): A patient-centered approach to grading evidence in the medical literature. *J Am Board Fam Pract* 2004; 17:59-67
60. Nyssen AS, Hansez I: Stress and burnout in anaesthesia. *Curr Opin Anaesthesiol* 2008; 21:406-11
61. Sharma A, Sharp DM, Walker LG, Monson JR: Stress and burnout in colorectal and vascular surgical consultants working in the UK National Health Service. *Psychooncology* 2008; 17:570-6 x'